

THE CLASSIFICATION OF TILTING MODULES OVER HARADA ALGEBRAS

KOTA YAMAURA

ABSTRACT. In the 1980s, Harada introduced a new class of algebras now called Harada algebras. Harada algebras provides us with a rich source of Auslander's 1-Gorenstein algebras. In this paper, we have two main results about Harada algebras. The first is the classification of modules over Harada algebras whose projective dimension is at most one. The second is the classification of tilting modules over Harada algebras, which is shown by giving a bijection between tilting modules over Harada algebras and tilting modules over direct products of upper triangular matrix algebras over K . A combinatorial description of tilting modules over upper triangular matrix algebras over K is known. These facts allow us to classify tilting modules over a given Harada algebra.

1. MAIN RESULTS

Two classes of algebras have been studied for a long time. The first is Nakayama algebras and the second is quasi-Frobenius algebras. In the 1980s, Harada introduced a new class of algebras now called Harada algebras, which give a common generalization of Nakayama algebras and quasi-Frobenius algebras. Many authors have studied the structure of Harada algebras (e.g. [7, 8, 17, 18, 19, 20, 21, 22]). Now let us recall that left Harada algebras are defined from a structural point of view as follows.

Definition 1.1. Let R be a basic algebra and $\text{Pi}(R)$ be a complete set of orthogonal primitive idempotents of R [1]. We call R a *left Harada algebra* if $\text{Pi}(R)$ can be arranged such that $\text{Pi}(R) = \{e_{ij}\}_{i=1}^m, j=1}^{n_i}$ where

- (1) $e_{i1}R$ is an injective R -module for any $i = 1, \dots, m$,
- (2) $e_{ij}R \simeq e_{i,j-1}J(R)$ for any $i = 1, \dots, m, j = 2, \dots, n_i$.

Here $J(R)$ is the Jacobson radical of R .

In [25], Thrall studied three properties of quasi-Frobenius algebras, called QF-1, QF-2, and QF-3. It follows from definition that left Harada algebras satisfy the property QF-3 which is the condition that the injective hull of the algebra is projective. This property is called 1-Gorenstein by Auslander (and dominant dimension at least one by Tachikawa) [5, 12, 14, 15, 24], and often plays an important role in the representation theory. Left Harada algebras form a class of 1-Gorenstein algebras, and their indecomposable projective modules have "nice" structure.

In this paper, we classify tilting modules over left Harada algebras. Tilting modules provide us a powerful tool in the representation theory of algebras and are due to [4, 9, 10].

Definition 1.2. Let R be an algebra. An R -module T is called a *partial tilting module* if T satisfies the following conditions.

- (1) $\text{proj.dim } T \leq 1$.
- (2) $\text{Ext}_R^1(T, T) = 0$.

Moreover, a partial tilting R -module T is called a *tilting module* if T satisfies the following condition.

- (3) There exists an exact sequence

$$0 \longrightarrow R_R \longrightarrow T_0 \longrightarrow T_1 \longrightarrow 0$$

where $T_0, T_1 \in \text{add}T$.

We can see from the above definition that tilting modules are a generalization of progenerators which appear in Morita's theorem. Morita's theorem shows that any progenerator P over an algebra R induces a category equivalence between $\text{mod}R$ and $\text{mod}(\text{End}_R(P))$. A generalization of Morita's theorem is the Brenner-Butler theorem. It says that any tilting module T over an algebra R induces two category equivalences between certain full subcategories of $\text{mod}R$ and of $\text{mod}(\text{End}_R(T))$. As a consequence of the Brenner-Butler theorem, some problems about R can be shifted to those of $\text{End}_R(T)$ (for example, finiteness of global dimension). By this reason, tilting modules are important for the study of algebras and a finding classification of tilting modules over a given algebra is an important problem in representation theory. The aim of this paper is to give a classification of tilting modules over a left Harada algebra.

Now we present the two main results of this paper. Let R be a left Harada algebra as in Definition 1.1. We denote by $J(M)$ the Jacobson radical of the R -module M , by $J^k(M)$ the k -th Jacobson radical of M and by $S(M)$ the socle of M . We put

$$(1.1) \quad P_{ij} := J^{j-1}(e_{i1}R) \simeq e_{ij}R \quad (1 \leq i \leq m, 1 \leq j \leq n_i)$$

for simplicity. Then we have a chain

$$P_{i1} \supset P_{i2} \supset \cdots \supset P_{in_i}$$

of indecomposable projective R -modules.

The first main result is the classification of R -modules whose projective dimension is equal to one. The full subcategory of the module category over a 1-Gorenstein algebra that consists of modules whose projective dimension is at most one is contravariantly finite ([14, 15]). Actually, we have a stronger result for left Harada algebra. Namely, the full subcategory of the module category over a left Harada algebra that consists of modules whose projective dimension is at most one is finite. It is obvious that $\text{proj.dim}(P_{ik}/P_{il}) = 1$. We can show converse.

Theorem 1.3. *A complete set of isomorphism classes of indecomposable R -modules whose projective dimension is equal to one is given as follows.*

$$\{P_{ik}/P_{il} \mid 1 \leq i \leq m, 1 \leq k < l \leq n_i\}.$$

This will be proved in Section 2 by using key lemmas which follow from Definition 1.1 directly.

The other main result is the classification of basic tilting R -modules. We denote by $\text{tilt}(R)$ the set of isomorphism classes of basic tilting R -modules and by $T_n(K)$ the $n \times n$ upper triangular matrix algebra over K

$$\begin{pmatrix} K & \cdots & K \\ & \ddots & \vdots \\ 0 & & K \end{pmatrix}.$$

The following asserts that tilting R -modules are described by tilting modules over a direct product of algebras of the form $T_n(K)$ which gives a typical example of a Harada algebra.

Theorem 1.4. *There exists a bijection*

$$\text{tilt}(R) \longrightarrow \text{tilt}(T_{n_1}(K)) \times \text{tilt}(T_{n_2}(K)) \times \cdots \times \text{tilt}(T_{n_m}(K)).$$

We will construct the above correspondence in Section 4. By the well-known classification of tilting modules over upper triangular matrix algebras over K which we recall in Section 5, we can classify tilting modules over a given left Harada algebra by the above correspondence.

Moreover, we can describe tilting $T_n(K)$ -modules combinatorially by using non-crossing partitions of a regular $(n+2)$ -polygon. In particular, we have the following application.

Corollary 1.5. *The number of basic tilting R -modules is equal to*

$$\prod_{i=1}^m \frac{1}{n_i + 1} \binom{2n_i}{n_i}.$$

Throughout this paper, an algebra means a finite dimensional associative algebra over an algebraically closed field K . We always deal with finitely generated right modules over algebras.

Acknowledgement The author is deeply grateful to Professor Kiyoichi Oshiro for giving me a chance to study ring theory and his warm encouragement. The author would like to thank Professor Osamu Iyama for generous support and suggestions getting to the points, and Martin Herschend and Michael Wemyss for helpful comments and suggestions.

2. PROOF OF THEOREM 1.3

In this section, we first give some key lemmas of this paper, that is, the properties of homomorphisms between indecomposable projective R -modules. Next we prove Theorem 1.3 by using these lemmas.

Let R be a left Harada algebra as in Definition 1.1. We use the notation (1.1).

Lemma 2.1. *If a submodule of P_{i1} is not contained in $J(P_{in_i})$, then it is P_{ij} for some $1 \leq j \leq n_i$.*

Proof. It follows from Definition 1.1 (b). $\square$

Lemma 2.2. *Let $f : P_{ij} \rightarrow P_{kl}$ be a homomorphism. Then the following hold.*

- (1) *f is monic if and only if $i = k$, $j \geq l$ and $\text{Im } f = P_{kj}$.*
- (2) *f is not monic if and only if $\text{Im } f \subset J(P_{kn_k})$.*
- (3) *Assume $i = k$ and $j < l$, we have $\text{Im } f \subset J(P_{kn_k})$*
- (4) *Assume $i \neq k$, we have $\text{Im } f \subset J(P_{kn_k})$.*

Proof. (1) We assume that f is monic. Then $i = k$ since $S(P_{ij}) \simeq S(P_{kl})$ and these are simple. By $\text{length}(P_{ij}) \leq \text{length}(P_{kl})$, we have $j \geq l$. By Lemma 2.1, the only submodule of P_{kl} whose length is equal to $\text{length}(P_{ij})$ is P_{kj} . The converse follows from $\text{length}(P_{ij}) = \text{Im } f$.

(2) We assume that $\text{Im } f \not\subset J(P_{kn_k})$. By Lemma 2.1, there exists $0 \leq r \leq n_k - l$ such that $\text{Im } f = P_{k,r+l}$. Therefore f is monic since f can be seen as an epimorphism between indecomposable projective R -modules. The converse follows from (1).

(3) Since $\text{length}(P_{ij}) > \text{length}(P_{il})$, there exists no monomorphism from P_{ij} to P_{il} . Therefore the assertion follows from (2).

(4) Since $i \neq k$, $S(P_{ij})$ and $S(P_{kl})$ are not isomorphic. Hence there exists no monomorphism from P_{ij} to P_{kl} . Therefore the assertion follows from (2). $\square$

Lemma 2.3. *Let $f : P_{ij} \rightarrow P_{il}$ be any monomorphism with $j \geq l$. Then the following hold.*

- (1) *For any homomorphism $g : P_{ij} \rightarrow P_{il'}$ with $l \geq l'$, there exists a homomorphism $h : P_{il} \rightarrow P_{il'}$ such that $g = hf$.*
- (2) *For any homomorphism $g : P_{ij} \rightarrow P_{st}$ which is not monic, there exists a homomorphism $h : P_{il} \rightarrow P_{st}$ such that $g = hf$.*
- (3) *For any homomorphism $g : P_{il'} \rightarrow P_{il}$ with $l' \geq j$, there exists a homomorphism $h : P_{il'} \rightarrow P_{ij}$ such that $g = fh$.*
- (4) *For any homomorphism $g : P_{st} \rightarrow P_{il}$ which is not monic, there exists a homomorphism $h : P_{st} \rightarrow P_{ij}$ such that $g = fh$.*

Proof. (1) Let $u : P_{il'} \rightarrow P_{i1}$ be the inclusion map. Since P_{i1} is injective, there exists a homomorphism $h : P_{il} \rightarrow P_{i1}$ such that $ug = hf$.

$$\begin{array}{ccccc}
 0 & \longrightarrow & P_{ij} & \xrightarrow{f} & P_{il} \\
 & & \downarrow g & & \searrow h \\
 & & P_{il'} & & \\
 & & \downarrow u & & \\
 & & P_{i1} & &
 \end{array}$$

Since $l \geq l'$, we have $\text{Im } h \subset P_{il'}$. We can see h as $h : P_{il} \rightarrow P_{il'}$.

(2) Let $u : P_{st} \rightarrow P_{s1}$ be the inclusion map. Since P_{s1} is injective, there exists a homomorphism $h : P_{il} \rightarrow P_{s1}$ such that $ug = hf$.

$$\begin{array}{ccccc}
 0 & \longrightarrow & P_{ij} & \xrightarrow{f} & P_{il} \\
 & & \downarrow g & & \searrow h \\
 & & P_{st} & & \\
 & & \downarrow u & & \\
 & & P_{s1} & &
 \end{array}$$

If h is monic, then $ug = hf$ is monic, hence g is monic. This is contradiction. Therefore h is not monic. By Lemma 2.2 (2), we have $\text{Im } h \subset J(P_{sn_s}) \subset P_{st}$. We can see h as $h : P_{il} \rightarrow P_{st}$.

(3) By Lemma 2.2 (1),(3), we have $\text{Im } f = P_{ij}$ and $\text{Im } g \subset P_{il'}$. Since $l' \geq j$, we have $\text{Im } g \subset \text{Im } f$. Since $P_{il'}$ is projective, there exists a homomorphism $h : P_{il'} \rightarrow P_{ij}$ such that $g = fh$.

$$\begin{array}{ccccc}
 & & P_{il'} & & \\
 & & \downarrow g & & \\
 & & P_{ij} & \xrightarrow{f} & P_{ij} \longrightarrow 0 \\
 & \nearrow h & & &
 \end{array}$$

(4) By Lemma 2.2 (1),(2), we have $\text{Im } g \subset J(P_{in_i}) \subset P_{ij} = \text{Im } f$. The assertion follows by the same argument as in the proof of (3). $\square$

The following result gives Theorem 1.3.

Lemma 2.4. *Let Q_i and Q'_j be indecomposable projective R -modules and*

$$f : Q := Q_1 \oplus \cdots \oplus Q_k \longrightarrow Q' := Q'_1 \oplus \cdots \oplus Q'_l$$

a monomorphism. Then there exists automorphisms $\varphi \in \text{Aut}_R(Q)$, $\psi \in \text{Aut}_R(Q')$ such that

$$\psi f \varphi^{-1} = \begin{pmatrix} f_1 & & 0 \\ & \ddots & \\ 0 & & f_k \\ \hline & 0 & \\ & \vdots & \\ & 0 & \end{pmatrix} : Q_1 \oplus \cdots \oplus Q_k \longrightarrow Q'_1 \oplus \cdots \oplus Q'_l.$$

Proof. We proceed by induction on k . First we consider the case $k = 1$. Then Q is an indecomposable projective R -module. We write f as

$$f : Q \xrightarrow{\begin{pmatrix} f_1 \\ \vdots \\ f_l \end{pmatrix}} Q', \quad f_i : Q \longrightarrow Q_i \quad (1 \leq i \leq l).$$

Since $S(Q)$ is simple, there exists an monomorphism in $\{f_1, \dots, f_l\}$. So we can assume that $f_1, \dots, f_r$ are monic and $f_{r+1}, \dots, f_l$ are not monic. We assume that $\text{length}(Q'_1) \leq \text{length}(Q'_i)$ for $2 \leq i \leq r$. Then for any $2 \leq j \leq r$ there exists a homomorphism $h_j : Q'_1 \longrightarrow Q'_j$ such that $f_j = h_j f_1$ by Lemma 2.3 (1). Moreover, for any $r+1 \leq j \leq n$ there exists a homomorphism $h_j : Q'_1 \longrightarrow Q'_j$ such that $f_j = h_j f_1$ by Lemma 2.3 (2). Let

$$\psi = \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ -h_2 & 1 & & 0 \\ \vdots & & \ddots & \\ -h_l & 0 & & 1 \end{array} \right) \in \text{Aut}_R(Q'_1 \oplus \cdots \oplus Q'_l).$$

Then we have

$$\psi f = \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ -h_2 & 1 & & 0 \\ \vdots & & \ddots & \\ -h_l & 0 & & 1 \end{array} \right) \begin{pmatrix} f_1 \\ \vdots \\ f_l \end{pmatrix} = \begin{pmatrix} f_1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}.$$

Next we assume that $k \geq 2$ and that the assertion holds for $k-1$. We assume that $\text{length}(Q_k) \leq \text{length}(Q_i)$ for $1 \leq i \leq k-1$. By applying the induction hypotheses to $f|_{Q_1 \oplus \cdots \oplus Q_{k-1}}$, we can assume that

$$f|_{Q_1 \oplus \cdots \oplus Q_{k-1}} : Q_1 \oplus \cdots \oplus Q_{k-1} \xrightarrow{\begin{pmatrix} f_1 & & 0 \\ & \ddots & \\ 0 & & f_{k-1} \\ \hline & & 0 \\ & & \vdots \\ & & 0 \end{pmatrix}} Q', \quad f_i : Q_i \longrightarrow Q'_i \quad (1 \leq i \leq k-1).$$

Therefore we can write f as

$$f : Q \xrightarrow{\begin{pmatrix} f_1 & & 0 & | & g_1 \\ & \ddots & & | & \vdots \\ 0 & & f_{k-1} & | & g_{k-1} \\ \hline 0 & \cdots & 0 & | & g_k \\ \vdots & & \vdots & | & \vdots \\ 0 & \cdots & 0 & | & g_l \end{pmatrix}} Q', \quad g_i : Q_k \longrightarrow Q'_i \quad (1 \leq i \leq l).$$

By Lemma 2.3 (3),(4), and the assumption on Q_k , for any $1 \leq i \leq k-1$ there exists a homomorphism $h_i : Q_k \longrightarrow Q_i$ such that $g_i = f_i h_i$. Let

$$\varphi = \left(\begin{array}{ccc|c} 1 & & 0 & h_1 \\ & \ddots & & \vdots \\ 0 & & 1 & h_{k-1} \\ \hline 0 & & & 1 \end{array} \right) \in \text{Aut}(Q_1 \oplus \cdots \oplus Q_k).$$

Then we have

$$\left(\begin{array}{ccc|c} f_1 & & 0 & \\ & \ddots & & \\ 0 & & f_{k-1} & 0 \\ \hline 0 & \cdots & 0 & g_k \\ \vdots & & \vdots & \vdots \\ 0 & \cdots & 0 & g_l \end{array} \right) \varphi = \left(\begin{array}{ccc|c} f_1 & & 0 & g_1 \\ & \ddots & & \vdots \\ 0 & & f_{k-1} & g_{k-1} \\ \hline 0 & \cdots & 0 & g_k \\ \vdots & & \vdots & \vdots \\ 0 & \cdots & 0 & g_l \end{array} \right) = f.$$

By applying the same argument as in the case $k = 1$ to

$$\begin{pmatrix} g_k \\ \vdots \\ g_l \end{pmatrix},$$

the assertion follows. $\square$

Now we can prove Theorem 1.3.

Proof. The projective dimension of P_{ik}/P_{il} is obviously equal to 1. Let X be an indecomposable R -module whose projective dimension is equal to one. Then there exists an exact sequence

$$0 \longrightarrow Q \longrightarrow Q' \longrightarrow X \longrightarrow 0$$

such that Q and Q' are projective R -modules. By Lemma 2.4 and since X is an indecomposable R -module, Q and Q' must be indecomposable R -modules. By Lemma 2.2 (1), X is isomorphic to one of P_{ik}/P_{il} . $\square$

3. TRIANGULAR FACTOR ALGEBRAS OF HARADA ALGEBRAS

In this section, we keep the notations from the previous section. We define a special factor algebra $\overline{R} = R/I$ of R which is isomorphic to a direct product $T_{n_1}(K) \times \cdots \times T_{n_m}(K)$ of upper triangular matrix algebras over K . The algebra $\overline{R}$ contains important information about R which is seen in Lemma 4.1 and Proposition 4.7.

We start by giving the following ideal I of R . We put

$$e_{ij}R \supset I_{ij} := J^{n_i-j+1}(e_{ij}R) \quad (1 \leq i \leq m, 1 \leq j \leq n_i),$$

and

$$R \supset I := \bigoplus_{i=1}^m \bigoplus_{j=1}^{n_i} I_{ij}.$$

Then we have the following result.

Lemma 3.1. *I is an ideal of R .*

Proof. Clearly I is a right ideal of R . We show I is a left ideal of R . It is enough to show that $rx \in I_{kl} = J^{n_k-l+1}(e_{kl}R)$ for any $x \in I_{ij} = J^{n_i-j+1}(e_{ij}R)$ and any $r \in e_{kl}R$. We consider the homomorphism

$$\varphi_r : I_{ij} \ni x \longmapsto rx \in e_{kl}R$$

of right R -modules. Since I_{ij} is indecomposable and non-projective, we have $\text{Im} \varphi_r \subset J^{n_k-l+1}(e_{kl}R) = I_{kl}$. Therefore I is a left ideal of R . $\square$

By Lemma 3.1, we can consider the factor algebra

$$\overline{R} := R/I.$$

We put

$$e_i := e_{i1} + e_{i2} + \cdots + e_{in_i}$$

for $1 \leq i \leq m$.

Now we show the following description of $\overline{R}$.

Proposition 3.2. *Under the hypotheses above, the following assertions hold.*

- (1) $\{e_i \mid 1 \leq i \leq m\}$ is a set of orthogonal central idempotents of $\overline{R}$ and there exists a K -algebra isomorphism

$$\overline{R}e_j \simeq T_{n_j}(K).$$

- (2) There exists a K -algebra isomorphism

$$\overline{R} \simeq T_{n_1}(K) \times T_{n_2}(K) \times \cdots \times T_{n_m}(K).$$

To prove the above proposition, we describe all indecomposable projective $\overline{R}$ -modules as factor modules of indecomposable projective R -modules. Since $I \subset J(R)$, we have that

$$\{e_{ij} + I \in \overline{R} \mid 1 \leq i \leq m, 1 \leq j \leq n_i\}$$

is a complete set of orthogonal primitive idempotents of $\overline{R}$. By the $\overline{R}$ -module isomorphism

$$e_{ij}\overline{R} = e_{ij}R/J^{n_i-j+1}(e_{ij}R) \simeq P_{ij}/J^{n_i-j+1}(P_{ij}) = P_{ij}/J(P_{in_i}),$$

a complete set of indecomposable projective $\overline{R}$ -modules is

$$\{P_{ij}/J(P_{in_i}) \mid 1 \leq i \leq m, 1 \leq j \leq n_i\}.$$

By Definition 1.1 (b),

$$(3.1) \quad 0 \subset P_{in_i}/J(P_{in_i}) \subset P_{i,n_i-1}/J(P_{in_i}) \subset \cdots \subset P_{i,j+1}/J(P_{in_i}) \subset P_{ij}/J(P_{in_i})$$

is a unique composition series of $P_{ij}/J(P_{in_i})$ as an $\overline{R}$ -module. Therefore any indecomposable projective $\overline{R}$ -module is serial and its composition factors are not isomorphic to each other.

From the above argument, we can prove Proposition 3.2.

Proof. (1) We calculate $\text{Hom}_{\overline{R}}(\overline{P}_{i,j}, \overline{P}_{k,l})$. If $i \neq k$, $\overline{P}_{i,j}$ and $\overline{P}_{k,l}$ have no common composition factors. So we have

$$\text{Hom}_{\overline{R}}(\overline{P}_{i,j}, \overline{P}_{k,l}) = 0.$$

If $i = k$, we can easily see that

$$\text{Hom}_{\overline{R}}(\overline{P}_{i,j}, \overline{P}_{i,l}) \simeq \begin{cases} K & (j \geq l) \\ 0 & (j < l) \end{cases}$$

by composition series (3.1).

Thus we have the following isomorphisms as K -vector space.

$$e_i \bar{R} e_j \simeq \text{Hom}_{\bar{R}}(e_j \bar{R}, e_i \bar{R}) \simeq \begin{pmatrix} \text{Hom}_{\bar{R}}(\bar{P}_{j,1}, \bar{P}_{i,1}) & \text{Hom}_{\bar{R}}(\bar{P}_{j,2}, \bar{P}_{i,1}) & \cdots & \text{Hom}_{\bar{R}}(\bar{P}_{j,n_j}, \bar{P}_{i,1}) \\ \text{Hom}_{\bar{R}}(\bar{P}_{j,1}, \bar{P}_{i,2}) & \text{Hom}_{\bar{R}}(\bar{P}_{j,2}, \bar{P}_{i,2}) & \cdots & \text{Hom}_{\bar{R}}(\bar{P}_{j,n_j}, \bar{P}_{i,2}) \\ \vdots & \vdots & \ddots & \vdots \\ \text{Hom}_{\bar{R}}(\bar{P}_{j,1}, \bar{P}_{i,n_i}) & \text{Hom}_{\bar{R}}(\bar{P}_{j,2}, \bar{P}_{i,n_i}) & \cdots & \text{Hom}_{\bar{R}}(\bar{P}_{j,n_j}, \bar{P}_{i,n_i}) \end{pmatrix}$$

$$\simeq \begin{cases} \begin{pmatrix} K & K & \cdots & K \\ & K & \cdots & K \\ & & \ddots & \vdots \\ 0 & & & K \end{pmatrix} & (i = j) \\ 0 & (i \neq j). \end{cases}$$

It is easily seen that the above isomorphism gives a K -algebra isomorphism when $i = j$.

(2) By (1), we have the following K -algebra isomorphism.

$$\bar{R} \simeq \begin{pmatrix} e_1 \bar{R} e_1 & e_1 \bar{R} e_2 & \cdots & e_1 \bar{R} e_m \\ e_2 \bar{R} e_1 & e_2 \bar{R} e_2 & \cdots & e_2 \bar{R} e_m \\ \vdots & \vdots & \ddots & \vdots \\ e_m \bar{R} e_1 & e_m \bar{R} e_2 & \cdots & e_m \bar{R} e_m \end{pmatrix} \simeq \begin{pmatrix} T_{n_1}(K) & & & 0 \\ & T_{n_2}(K) & & \\ & & \ddots & \\ 0 & & & T_{n_m}(K) \end{pmatrix}.$$

□

4. PROOF OF THEOREM 1.4

In this section, we keep the notations from the previous section. The aim of this section is to prove Theorem 1.4. First we describe the indecomposable $\bar{R}$ -modules by using the indecomposable projective R -modules P_{ij} . Next we consider a certain functor F preserving the vanishing property $\text{Ext}^1(-, -) = 0$, from the category $\mathcal{P}$ of R -modules whose projective dimension is at most one to $\text{mod } \bar{R}$. We construct a bijection between $\text{tilt}(R)$ and $\text{tilt}(\bar{R})$ by using the functor F which gives Theorem 1.4.

We start by giving a classification of indecomposable $\bar{R}$ -modules. By Proposition 3.2, it is shown that $\bar{R}$ is a Nakayama algebra. It is well-known that any indecomposable module over a Nakayama algebra is isomorphic to some subfactor of an indecomposable projective module ([3]). By this fact and the unique composition series (3.1) of $P_{ij}/J(P_{in_i})$,

$$\{P_{ik}/P_{il} \mid 1 \leq i \leq m, 1 \leq k < l \leq n_i\}$$

is a complete set of indecomposable nonprojective $\bar{R}$ -modules.

We put

$$\bar{P}_{i,j} := P_{ij}/J(P_{in_i}) \simeq e_{ij} \bar{R}$$

for $1 \leq i \leq m, 1 \leq j \leq n_i$ and

$$P_{i,k,l} := P_{ik}/P_{il}$$

for $1 \leq i \leq m, 1 \leq k < l \leq n_i$ for simplicity. These R -modules can be regarded as $\bar{R}$ -modules.

We have the following diagram for any $1 \leq i \leq m$ by our definitions.

$$\begin{array}{ccccccc}
\overline{P}_{i,1} & \longrightarrow & P_{i,1,n_i} & \longrightarrow & P_{i,1,n_i-1} & \longrightarrow & \cdots \longrightarrow P_{i,1,3} \longrightarrow P_{i,1,2} \\
\cup & & \cup & & \cup & & \cup \\
\overline{P}_{i,2} & \longrightarrow & P_{i,2,n_i} & \longrightarrow & P_{i,2,n_i-1} & \longrightarrow & \cdots \longrightarrow P_{i,2,3} \\
\cup & & \cup & & \cup & & \\
\vdots & & \vdots & & \vdots & & \\
\cup & & \cup & & \cup & & \\
\overline{P}_{i,n_i-2} & \longrightarrow & P_{i,n_i-2,n_i} & \longrightarrow & P_{i,n_i-2,n_i-1} & & \\
\cup & & \cup & & & & \\
\overline{P}_{i,n_i-1} & \longrightarrow & P_{i,n_i-1,n_i} & & & & \\
\cup & & & & & & \\
\overline{P}_{i,n_i} & & & & & &
\end{array}$$

In the above diagram, right arrows mean natural epimorphisms. We remark that the above diagram is the AR-quiver of $\text{mod}(\overline{Re}_i)$ (see Section 5).

Let $\mathcal{P}$ be the category of R -modules whose projective dimension is at most one. We define the full subcategories $\mathcal{P}_i$ of $\mathcal{P}$ for $1 \leq i \leq m$ by

$$\mathcal{P}_i := \text{add}\{P_{ij}, P_{i,k,l} \mid 1 \leq j \leq n_i, 1 \leq k < l \leq n_i\}.$$

By Theorem 1.3, we have

$$\mathcal{P} = \text{add}(\mathcal{P}_1 \cup \mathcal{P}_2 \cup \cdots \cup \mathcal{P}_m).$$

A key role is played by the functor

$$F := - \otimes_R \overline{R} : \mathcal{P} \longrightarrow \text{mod} \overline{R}.$$

Lemma 4.1. *Under the hypotheses above, the following hold.*

- (1) *The functor F induces a bijection between the isomorphism classes of R -modules which lie in $\mathcal{P}$ and the isomorphism classes of $\overline{R}$ -modules.*
- (2) *The restriction on F to $\mathcal{P}_i$ induces a one to one correspondence between the isomorphism classes of $\mathcal{P}_i$ and the isomorphism classes of $\text{mod}(\overline{Re}_i)$.*

Proof. We calculate $F(M)$ for an indecomposable R -module M which lies in $\mathcal{P}$. We have isomorphisms

$$F(P_{ij}) = P_{ij} \otimes_R \overline{R} \simeq P_{ij}/(P_{ij}I) = P_{ij}/J^{n_i}(P_{i1}) = \overline{P}_{ij},$$

$$F(P_{i,k,l}) = P_{i,k,l} \otimes_R \overline{R} \simeq P_{i,k,l}/(P_{i,k,l}I) = P_{i,k,l}.$$

for $1 \leq i \leq m$, $1 \leq j \leq n_i$ and $1 \leq k < l \leq n_i$. The assertion follows. $\square$

Now we state a theorem that gives a bijection between $\text{tilt}(R)$ and $\text{tilt}(\overline{R})$ by using the functor F .

Theorem 4.2. *Under the hypotheses above, every tilting R -module lies in $\mathcal{P}$ and we have a bijection*

$$F : \text{tilt}(R) \ni T \longmapsto F(T) \in \text{tilt}(\overline{R}).$$

As a consequence of Theorem 4.2, we have the following corollary immediately.

Corollary 4.3. *Under the hypotheses above, we have a bijection*

$$\text{tilt}(R) \ni T \longmapsto (F(T)e_1, \dots, F(T)e_m) \in \text{tilt}(\overline{Re}_1) \times \cdots \times \text{tilt}(\overline{Re}_m).$$

Hence by Proposition 3.2, we have Theorem 1.4. In the rest of this section, we prove of Theorem 4.2 .

We have to know when $\text{Ext}_R^1(X, Y) = 0$ holds for $X, Y \in \mathcal{P}$. We start with the following result for $\text{Ext}_R^1(\mathcal{P}_i, \mathcal{P}_u)$.

Lemma 4.4. $\text{Ext}_R^1(\mathcal{P}_i, \mathcal{P}_u) = 0$ if $i \neq u$.

Proof. It is obvious that $\text{Ext}_R^1(\mathcal{P}_{ij}, \mathcal{P}_u) = 0$. We show $\text{Ext}_R^1(\mathcal{P}_{i,k,l}, \mathcal{P}_u) = 0$ for $1 \leq k < l \leq n_i$.

First we show $\text{Ext}_R^1(\mathcal{P}_{i,k,l}, P_{uv}) = 0$ for $1 \leq v \leq n_u$. We take a projective resolution

$$(4.1) \quad 0 \longrightarrow P_{il} \xrightarrow{f} P_{ik} \longrightarrow P_{i,k,l} \longrightarrow 0$$

of $P_{i,k,l}$ in $\text{mod} R$. By applying $\text{Hom}_R(-, P_{uv})$ to the above exact sequence, we have an exact sequence

$$\text{Hom}_R(P_{ik}, P_{uv}) \xrightarrow{\text{Hom}(f, P_{uv})} \text{Hom}_R(P_{il}, P_{uv}) \longrightarrow \text{Ext}_R^1(P_{i,k,l}, P_{uv}) \longrightarrow 0.$$

By the assumption $i \neq u$, there is no monomorphism from P_{il} to P_{uv} since the simple socles $S(P_{il})$ and $S(P_{uv})$ are not isomorphic. By Lemma 2.3 (2), $\text{Hom}(f, P_{uv})$ is an epimorphism. Therefore we have $\text{Ext}_R^1(P_{i,k,l}, P_{uv}) = 0$.

Next we show $\text{Ext}_R^1(P_{i,k,l}, P_{u,s,t}) = 0$ for $1 \leq s < t \leq n_u$. By applying $\text{Hom}_R(-, P_{u,s,t})$ to (4.1), we have an exact sequence

$$\text{Hom}_R(P_{ik}, P_{u,s,t}) \longrightarrow \text{Hom}_R(P_{il}, P_{u,s,t}) \longrightarrow \text{Ext}_R^1(P_{i,k,l}, P_{u,s,t}) \longrightarrow 0.$$

By the assumption $i \neq u$, $P_{il}/J(P_{il})$ does not appear in composition factors of $P_{u,s,t}$. Therefore we have $\text{Hom}_R(P_{il}, P_{u,s,t}) = 0$, and so $\text{Ext}_R^1(P_{i,k,l}, P_{u,s,t}) = 0$. $\square$

Next we consider $\text{Ext}_R^1(\mathcal{P}_i, \mathcal{P}_i) = 0$. We need the following result.

Lemma 4.5. For any $1 \leq i \leq m$, $1 \leq j \leq n_i$ and $1 \leq k < l \leq n_i$, the natural epimorphism $\varphi : P_{ij} \longrightarrow \overline{P}_{i,j}$ induces an isomorphism

$$\text{Hom}(\varphi, P_{i,k,l}) : \text{Hom}_{\overline{R}}(\overline{P}_{i,j}, P_{i,k,l}) = \text{Hom}_R(\overline{P}_{i,j}, P_{i,k,l}) \longrightarrow \text{Hom}_R(P_{ij}, P_{i,k,l}).$$

Proof. It is obvious that $\text{Hom}_R(\overline{P}_{i,j}, P_{i,k,l}) = \text{Hom}_{\overline{R}}(\overline{P}_{i,j}, P_{i,k,l})$ holds. We show that $\text{Hom}(\varphi, P_{i,k,l})$ is an isomorphism.

Since φ is epic, we have that $\text{Hom}(\varphi, P_{i,k,l})$ is monic. Since any $f \in \text{Hom}_R(P_{ij}, P_{i,k,l})$ satisfies $\text{Ker } f \supset P_{ij}I = \text{Ker } \varphi$, we have that f factors through φ . Thus $\text{Hom}(\varphi, P_{i,k,l})$ is an isomorphism. $\square$

Proposition 4.6. The following hold.

- (1) For $1 \leq j \leq n_i$, $\text{Ext}_R^1(P_{ij}, \mathcal{P}) = 0 = \text{Ext}_R^1(\overline{P}_{i,j}, \text{mod } \overline{R})$.
- (2) For $1 \leq k < l \leq n_i$, $1 \leq s < t \leq n_i$, there is a K -vector space isomorphism

$$\text{Ext}_R^1(P_{i,k,l}, P_{i,s,t}) \simeq \text{Ext}_{\overline{R}}^1(P_{i,k,l}, P_{i,s,t}).$$

- (3) For $1 \leq k < l \leq n_i$, $1 \leq j \leq n_i$, $\text{Ext}_R^1(P_{i,k,l}, P_{ij}) = 0$ if and only if $\text{Ext}_{\overline{R}}^1(P_{i,k,l}, \overline{P}_{i,j}) = 0$.

Proof. (1) Obvious.

(2) We have a natural projective resolution

$$(4.2) \quad 0 \longrightarrow P_{il} \xrightarrow{f} P_{ik} \longrightarrow P_{i,k,l} \longrightarrow 0$$

of $P_{i,k,l}$ in $\text{mod} R$ and a natural projective resolution

$$(4.3) \quad 0 \longrightarrow \overline{P}_{i,l} \xrightarrow{f'} \overline{P}_{i,k} \longrightarrow P_{i,k,l} \longrightarrow 0$$

of $P_{i,k,l}$ in $\text{mod } \overline{R}$. For natural epimorphisms $\varphi : P_{ik} \longrightarrow \overline{P}_{i,k}$ and $\varphi' : P_{il} \longrightarrow \overline{P}_{i,l}$, we have the following commutative diagram.

$$\begin{array}{ccccccc} 0 & \longrightarrow & P_{il} & \xrightarrow{f} & P_{ik} & \longrightarrow & P_{i,k,l} \longrightarrow 0 \\ & & \downarrow \varphi' & & \downarrow \varphi & & \\ 0 & \longrightarrow & \overline{P}_{i,l} & \xrightarrow{f'} & \overline{P}_{i,k} & \longrightarrow & P_{i,k,l} \longrightarrow 0 \end{array}$$

By applying $\text{Hom}_R(-, P_{i,s,t})$ to the upper row and applying $\text{Hom}_{\overline{R}}(-, P_{i,s,t})$ to the lower row, we have the following commutative diagram.

$$\begin{array}{ccccc} \text{Hom}_{\overline{R}}(\overline{P}_{i,k}, P_{i,s,t}) & \longrightarrow & \text{Hom}_{\overline{R}}(\overline{P}_{i,l}, P_{i,s,t}) & \longrightarrow & \text{Ext}_{\overline{R}}^1(P_{i,k,l}, P_{i,s,t}) \longrightarrow 0 \\ \parallel & & \parallel & & \\ \text{Hom}_R(\overline{P}_{i,k}, P_{i,s,t}) & & \text{Hom}_R(\overline{P}_{i,l}, P_{i,s,t}) & & \\ \downarrow \text{Hom}(\varphi, P_{i,s,t}) & & \downarrow \text{Hom}(\varphi', P_{i,s,t}) & & \\ \text{Hom}_R(P_{ik}, P_{i,s,t}) & \longrightarrow & \text{Hom}_R(P_{il}, P_{i,s,t}) & \longrightarrow & \text{Ext}_R^1(P_{i,k,l}, P_{i,s,t}) \longrightarrow 0 \end{array}$$

By Lemma 4.5, $\text{Hom}(\varphi, P_{i,s,t})$ and $\text{Hom}(\varphi', P_{i,s,t})$ are isomorphisms. Consequently we have an isomorphism $\text{Ext}_R^1(P_{i,k,l}, P_{i,s,t}) \simeq \text{Ext}_{\overline{R}}^1(P_{i,k,l}, P_{i,s,t})$.

(3) By applying $\text{Hom}_R(-, P_{ij})$ to (4.2), we have an exact sequence

$$\text{Hom}_R(P_{ik}, P_{ij}) \xrightarrow{\text{Hom}(f, P_{ij})} \text{Hom}_R(P_{il}, P_{ij}) \longrightarrow \text{Ext}_R^1(P_{i,k,l}, P_{ij}) \longrightarrow 0.$$

It can be seen that $\text{Ext}_R^1(P_{i,k,l}, P_{ij}) = 0$ if and only if $\text{Hom}(f, P_{ij})$ is an epimorphism.

We show that $\text{Hom}(f, P_{ij})$ is an epimorphism if and only if $j \leq k$ or $l < j$. First we assume that $j > k$ and $l \geq j$. By $l \geq j$, there exists a monomorphism from P_{il} to P_{ij} . But there are no monomorphisms from P_{ik} to P_{ij} by $j > k$. Since P_{ik} has simple socle, gf is not monic for any $g \in \text{Hom}_R(P_{ik}, P_{ij})$. Thus $\text{Hom}(f, P_{ij})$ is not an epimorphism. Next we assume $j \leq k$. By Lemma 2.3 (1), $\text{Hom}(f, P_{ij})$ is an epimorphism. Finally we assume $l < j$. Then by $\text{length}(P_{il}) > \text{length}(P_{ij})$, there are no monomorphisms from P_{il} to P_{ij} . By Lemma 2.3 (2), $\text{Hom}(f, P_{ij})$ is an epimorphism.

On the other hand, by applying $\text{Hom}_{\overline{R}}(-, \overline{P}_{i,j})$ to (4.3), we have an exact sequence

$$\text{Hom}_{\overline{R}}(\overline{P}_{i,k}, \overline{P}_{i,j}) \xrightarrow{\text{Hom}(f', \overline{P}_{i,j})} \text{Hom}_{\overline{R}}(\overline{P}_{i,l}, \overline{P}_{i,j}) \longrightarrow \text{Ext}_{\overline{R}}^1(P_{i,k,l}, \overline{P}_{i,j}) \longrightarrow 0.$$

It can be seen that $\text{Ext}_{\overline{R}}^1(P_{i,k,l}, \overline{P}_{i,j}) = 0$ if and only if $\text{Hom}(f', \overline{P}_{i,j})$ is an epimorphism. We can show that $\text{Hom}(f', \overline{P}_{i,j})$ is an epimorphism if and only if $j \leq k$ or $l < j$ by the same argument as above.

Consequently $\text{Ext}_R^1(P_{i,k,l}, P_{ij}) = 0$ if and only if $j \leq k$ or $l < j$ which is equivalent to $\text{Ext}_{\overline{R}}^1(P_{i,k,l}, \overline{P}_{i,j}) = 0$. $\square$

By the Proposition 4.6, we have the following proposition.

Proposition 4.7. *For any $T \in \mathcal{P}$, we have that $\text{Ext}_R^1(T, T) = 0$ if and only if $\text{Ext}_{\overline{R}}^1(F(T), F(T)) = 0$.*

Proof. The assertion follows from Lemma 4.4 and Proposition 4.6. $\square$

We need the following well-known proposition which describes a very useful equivalent condition of tilting modules.

Proposition 4.8 ([3]). *Let R be an algebra and T a partial tilting R -module. Then the following are equivalent.*

- (1) *T is a tilting module.*
- (2) *The number of pairwise nonisomorphic indecomposable direct summand of T is equal to the number of pairwise nonisomorphic simple R -modules.*

Now we can prove Theorem 4.2.


Proof. First we take a basic tilting R -module T . Then T lies in $\mathcal{P}$ by Theorem 1.3. So we can consider $F(T)$. By Lemma 4.1, we have that F gives a bijection between isomorphism classes of R -modules whose projective dimension is at most one and those of $\overline{R}$ -modules. By Proposition 4.8 and the fact that the number of isomorphism classes of simple R -modules is equal to that of $\overline{R}$ -modules, we only have to show that $T \in \mathcal{P}$ satisfies $\text{Ext}_R^1(T, T) = 0$ if and only if $\text{Ext}_{\overline{R}}^1(F(T), F(T)) = 0$. This follows from Proposition 4.7. $\square$

5. COMBINATORIAL DESCRIPTION OF TILTING $T_n(K)$ -MODULES

First we recall the classification of basic tilting modules over the upper triangular matrix algebra $T_n(K)$. Our classification should be well-known for experts [2, 11, 16, 23]. Nevertheless we will give a complete proof since there does not seem to exist proper reference. Next we explain how to classify basic tilting modules over a given left Harada algebra.

We show the well-known classification of basic tilting $T_n(K)$ -modules by constructing a bijection between $\text{tilt}(T_n(K))$ and the set of non-crossing partitions of the regular $n + 2$ -polygon into triangles.


First we introduce coordinates in the AR-quiver of $T_n(K)$ as follows.



We remark that the vertex (i, j) corresponds the $T_n(K)$ -module

$$M_{ij} = (0 \cdots 0 \overset{j-2}{\underbrace{K}} \cdots \overset{1}{\underbrace{K}}) / (0 \cdots 0 \overset{i}{\underbrace{K}} \cdots \overset{1}{\underbrace{K}}) = (0 \cdots 0 \overset{j-2}{\underbrace{K}} \cdots \overset{i}{\underbrace{K}} \cdots \overset{1}{\underbrace{K}} 0 \cdots 0) .$$

Next we consider a regular $(n+2)$ -polygon R_{n+2} whose vertices are numbered as follows.



We denote by $D(R_{n+2})$ the set of all diagonals of R_{n+2} except edges of R_{n+2} . We call a subset S of $D(R_{n+2})$ a *non-crossing partition* of R_{n+2} if S satisfies the following conditions.

- (1) Any two distinct diagonals in S do not cross except at their endpoints.
- (2) R_{n+2} is divided into triangles by diagonals in S .

We denote by $\mathcal{P}_{n+2}$ the set of an non-crossing partitions of R_{n+2} .

Now we construct the correspondence Φ from $\mathcal{P}_{n+2}$ to $\text{tilt}(\text{T}_n(K))$. We take $S \in \mathcal{P}_{n+2}$. We remark that non-crossing partition of R_{n+2} consists of $n-1$ diagonals. We denote by (i, j) the diagonal between i and j for $i < j$ and put

$$S = \{(i_1, j_1), (i_2, j_2), \dots, (i_{n-1}, j_{n-1})\}.$$

Then we define

$$\Phi(S) := M_{1, n+2} \oplus \left(\bigoplus_{k=1}^{n-1} M_{i_k, j_k} \right).$$

It is shown that this is a basic tilting $\text{T}_n(K)$ -module.

Then the following hold.

Theorem 5.1. *The above correspondence Φ is a bijection.*

Proof. We divide the proof into five parts.

- (i) One can easily check that the following conditions are equivalent for $(i, j) \neq (i', j')$.

- (a) $\text{Ext}_R^1(M_{i,j}, M_{i',j'}) = 0$ and $\text{Ext}_R^1(M_{i',j'}, M_{i,j}) = 0$.
- (b) The diagonals (i, j) and (i', j') do not cross except at their endpoints.

- (ii) For any (i, j) , we have that $\text{Ext}_R^1(M_{i,j}, M_{i,j}) = 0$.

- (iii) $M_{1, n+2}$ is a projective injective $\text{T}_n(K)$ -module.


- (iv) If S is a non-crossing partition, then $\Phi(S)$ is a partial tilting module by (i), (ii) and (iii).

Since $\Phi(S)$ has n non-isomorphic indecomposable summands, it is a tilting module by Proposition 4.8.

- (v) Any basic tilting $\text{T}_n(K)$ -module T has $M_{1, n+2}$ as a summand. Since T has exactly n indecomposable direct summands, there exists a subset S of $D(R_{n+2})$ consists of $(n-1)$ elements such that $T = \Phi(S)$. By (i), S is non-crossing partition of R_{n+2} . $\square$

Theorem 5.1 gives a constructive bijection.

Example 5.2. We consider $n = 3$ case. We classify basic tilting $T_3(K)$ -modules by using Theorem 5.1. The partitions of the regular pentagon into triangles are given as follows.



Therefore the number of basic tilting $T_3(K)$ -modules is equal to 5 and all of basic tilting $T_3(K)$ -modules are given as follows.

- (1) $(K \ K \ K) \oplus (0 \ K \ K) \oplus (0 \ 0 \ K)$,
- (2) $(K \ K \ K) \oplus (K \ K \ 0) \oplus (0 \ K \ 0)$,
- (3) $(K \ K \ K) \oplus (K \ 0 \ 0) \oplus (0 \ 0 \ K)$,
- (4) $(K \ K \ K) \oplus (0 \ K \ K) \oplus (0 \ K \ 0)$,
- (5) $(K \ K \ K) \oplus (K \ K \ 0) \oplus (K \ 0 \ 0)$.

Now we show examples of classifications of tilting modules over left Harada algebras.

Example 5.3.

- (1) Let R be a local quasi-Frobenius algebra. Then we consider block extension (c.f. [8, 22])

$$R(n) = \begin{pmatrix} R & \cdots & R \\ & \ddots & \vdots \\ J(R) & & R \end{pmatrix}$$

for $n \in \mathbb{N}$ of R which is a subalgebra of $n \times n$ full matrix algebra over R . We can show that

- (a) the first row is a injective module,
- (b) the i -th row is the Jacobson radical of the $(i - 1)$ -th row for $2 \leq i \leq n$.

In particular $R(n)$ is a left Harada algebra with $m = 1$ and $n_1 = n$ in Definition 1.1.

By Corollary 4.3, we have a bijection $F : \text{tilt}(R(n)) \longrightarrow \text{tilt}(T_n(K))$. We can obtain all basic tilting $R(n)$ -modules from the definition of F and Theorem 5.1.

- (2) Let R be a basic quasi-Frobenius algebra which has complete set of orthogonal primitive idempotents $\{e, f\}$. Then we can represent R as follows (c.f. [1]).

$$R \simeq \begin{pmatrix} eRe & eRf \\ fRe & fRf \end{pmatrix}.$$

We put $Q := eRe$, $W := fRf$, $A := eRf$ and $B := fRe$. Now we consider the block extension (c.f. [8, 22])

$$R(n_1, n_2) = \left(\begin{array}{ccc|ccc} Q & \cdots & Q & A & \cdots & A \\ & & \vdots & \vdots & & \vdots \\ & & \ddots & \vdots & & \vdots \\ J(Q) & & Q & A & \cdots & A \\ \hline B & \cdots & B & W & \cdots & W \\ \vdots & & \vdots & & \ddots & \vdots \\ B & \cdots & B & J(W) & & W \end{array} \right)$$

for $n_1, n_2 \in \mathbb{N}$ of R which is a subalgebra of $\text{End}_R((eR)^{n_1} \oplus (fR)^{n_2})$. We can show that

- (a) the first and $n_1 + 1$ row are injective modules,
- (b) the i -th row is the Jacobson radical of the $(i - 1)$ -th row for $2 \leq i \leq n$ and $n + 2 \leq i \leq n + m$.

In particular $R(n_1, n_2)$ is a left Harada algebra with $m = 2$ in Definition 1.1.

By Corollary 4.3, we have a bijection $F : \text{tilt}(R(n_1, n_2)) \longrightarrow \text{tilt}(T_{n_1}(K)) \times \text{tilt}(T_{n_2}(K))$. We can obtain all basic tilting $R(n_1, n_2)$ -modules from the definition of F and Theorem 5.1.

REFERENCES

- [1] F. W. Anderson, K. R. Fuller: Rings and Categories of Modules (second edition), Graduate Texts in Math. 13, Springer-Verlag, Heidelberg/New York/Berlin (1991)
- [2] T. Araya: Exceptional Sequences over path algebras of type A_n and Non-crossing Spanning Trees, arXiv: 0904. 2831.
- [3] I. Assem, D. Simson, A. Skowronski: Elements of the Representation Theory of Associative Algebras, London Mathematical Society Student Texts 65, Cambridge university press (2006)
- [4] M. Auslander, M. I. Platzeck, I. Reiten: Coxeter functors without diagrams. Trans. Amer. Math. Soc. 250 (1979), 1-46
- [5] M. Auslander, I. Reiten: k -Gorenstein algebras and syzygy modules, J. Pure Appl. Algebra 92 (1994), 1-27.
- [6] M. Auslander, I. Reiten, S. Smalø: Representation Theory of Artin Algebras, Cambridge Studies in Advanced Mathematics 36, Cambridge university press (1995)
- [7] Y. Baba and K. Iwase: On quasi-Harada rings, J. Algebra 155 (1996), 415-434
- [8] Y. Baba and K. Oshiro: Classical Artinian Rings and Related Topics, Lecture note, preprint
- [9] S. Brenner, M. C. R. Butler: Generalizations of the Bernstein Gelfand Ponomarev reflection functors. Representation theory, II (Proc. Second Internat. Conf., Carleton Univ., Ottawa, Ont., 1979), Lecture Notes in Math., 832, Springer, Berlin-New York (1980), 103-169
- [10] I. N. Bernstein, I. M. Gelfand, V. A. Ponomarev: Coxeter functors, and Gabriel's theorem. (Russian) Uspehi Mat. Nauk 28 no. 2(170) (1973), 19-33
- [11] P. Caldero, F. Chapoton, R. Schiffler: Quivers with relations arising from clusters (A_n case), Trans. Amer. Math. Soc. 358 no. 3 (2006), 1347-1364
- [12] R. Fossum, P. Griffith, I. Reiten: Trivial extensions of abelian categories, Lecture Notes in Mathematics Vol. 456. Springer-Verlag, Berlin-New York, (1975)
- [13] M. Harada: Non-small modules and non-cosmall modules, Ring Theory. Proceedings of 1978 Antwerp Conference, New York (1979), 669-690
- [14] Z. Huang, O. Iyama: Auslander-type conditions and cotorsion pairs. J. Algebra 318 no. 1 (2007), 93-100
- [15] K. Igusa, S. O. Smalø, G. Todorov: Finite projectivity and contravariant finiteness. Proc. Amer. Math. Soc. 109 no. 4 (1990), 937-941
- [16] O. Iyama: Higher-dimensional Auslander-Reiten theory on maximal orthogonal subcategories, Adv. Math. 210 no. 1 (2007), 22-50
- [17] K. Koike: Almost self-duality and Harada rings, J. Algebra 254 (2002), 336-361
- [18] K. Oshiro: Lifting modules, extending modules and their applications to QF-rings, Hokkaido Math. J. 13 (1984), 310-338
- [19] K. Oshiro: Lifting modules, extending modules and their applications to generalized uniserial rings, Hokkaido Math. J. 13 (1984), 339-346
- [20] K. Oshiro: On Harada rings I, Math. J. Okayama Univ. 31 (1989), 161-178
- [21] K. Oshiro: On Harada rings II, Math. J. Okayama Univ. 31 (1989), 169-188
- [22] K. Oshiro: On Harada rings III, Math. J. Okayama Univ. 32 (1990), 111-118
- [23] C. Riedtmann: Representation-finite self-injective algebras of class A_n , Representation theory, II (Proc. Second Internat. Conf., Carleton Univ., Ottawa, Ont., 1979), Lecture Notes in Math., 832, Springer, Berlin (1980), 449-520
- [24] H. Tachikawa: Quasi-Frobenius rings and generalizations, Lecture Notes in Mathematics Vol. 351. Springer-Verlag, Berlin-New York, (1973)
- [25] Thrall, R. M.: Some generalization of quasi-Frobenius algebras, Trans. Amer. Math. Soc. 64 (1948), 173-183

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, FROCHO, CHIKUSAKU, NAGOYA, 464-8602, JAPAN

E-mail address: m07052d@math.nagoya-u.ac.jp